

Citation for published version:

Morgan, BJ & Madden, PA 2014, 'Molecular dynamics simulation of ionic transport at coherent interfaces in fluorite heterostructures', *Physical Review B*, vol. 89, no. 5, 054304.
<https://doi.org/10.1103/PhysRevB.89.054304>

DOI:

[10.1103/PhysRevB.89.054304](https://doi.org/10.1103/PhysRevB.89.054304)

Publication date:

2014

Document Version

Peer reviewed version

[Link to publication](#)

This is supplementary information to the authors' accepted version of an article published by the American Physical Society (APS) and available via: <http://dx.doi.org/10.1103/PhysRevB.89.054304>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Molecular Dynamics Simulation of Coherent Interfaces in Fluorite Heterostructures: Supplementary Information

Benjamin J. Morgan and Paul A. Madden

*Department of Materials, University of Oxford, Parks Road, Oxford OX1 3PH, UK**




FIG. S1: Defect formation energies for single vacancies and interstitials as a function of x for $\sigma_F = 0.04, 0.05, 0.06, 0.07$.

Fig. S1 plots relative energies as a function of σ_F for independent vacancies (upper panel) and interstitials (lower panel) at all possible x -positions in the simulation cell with fully relaxed geometries.

* benjamin.morgan@materials.ox.ac.uk